

# Preparation of Axially Chiral 2,2'-Biimidazole Ligands through Remote Chirality Delivery and Their Application in Asymmetric Carbene Insertion into N–H of Carbazoles

Hong-Qiang Shen,<sup>†,‡</sup> Bo Wu,<sup>†</sup> Huan-Ping Xie,<sup>†</sup> and Yong-Gui Zhou<sup>\*,†,§</sup>


<sup>†</sup>State Key Laboratory of Catalysis, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian 116023, People's Republic of China

<sup>‡</sup>University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China

<sup>§</sup>Collaborative Innovation Centre of Chemical Science and Engineering, Tianjin 300071, People's Republic of China

## Supporting Information


**ABSTRACT:** Axially chiral biimidazole ligands have been rarely synthesized and studied, in contrast to the significant achievements in the synthesis and application of central chiral imidazole ligands. Herein, a series of novel axially chiral 2,2'-biimidazole ligands were synthesized from the reaction of 2,2'-bis(bromomethyl)-1,1'-binaphthalene and 2,2'-biimidazole in one step through the strategy of remote chirality delivery. These ligands have been proven to be efficient for Cu- or Fe-catalyzed asymmetric insertion of  $\alpha$ -aryl- $\alpha$ -diazoacetates into the N–H bond of carbazoles with up to 96% ee.




Imidazoles are ubiquitous and prevalent structural motifs in biologically active molecules and bioprocesses.<sup>1</sup> It is well-known that imidazole served as the side chain of histidine can coordinate with metal ions in metalloproteins. Moreover, the imidazole side group of histidine has a  $pK_a$  of  $\sim 6$ , which exhibits amphoteric properties playing an active role in proton processes in biological systems (Scheme 1a).<sup>1d</sup> Inspired by its diverse biological properties, imidazoles have been designed as chiral organocatalysts (NMI, DPI, etc.)<sup>2</sup> and ligands by organic

## Scheme 1. Chiral Imidazole Compounds and the Design of Axially Chiral 2,2'-Biimidazole Ligands


### a) Chiral imidazole compound



### b) Representative Chiral Imidazole Organocatalyst and Ligands



### c) This Work: Axially Chiral 2,2'-Biimidazole Ligands



chemists for asymmetric catalysis in organic transformations. Especially, considering the excellent coordination ability of imidazoles with metal ions, the imidazole-based ligands have attracted considerable attention.<sup>3</sup> Many chiral ligands bearing imidazole moieties have been developed and applied in asymmetric hydrogenation, dehydrative allylation, aldol reaction, cycloaddition and Henry reaction, etc.<sup>4</sup> As the representative examples in this area, a series of imidazole-based N/P-ligand have been prepared and utilized in Ir-catalyzed asymmetric hydrogenation by Andersson's group in 2008.<sup>5</sup> Subsequently, Aponick's group successfully synthesized an imidazole-based biaryl N/P-ligand named StackPhos which atropisomer interconversion was inhibited by  $\pi$ - $\pi$  stacking. This ligand performed well in the asymmetric Cu-catalyzed A3-coupling reaction.<sup>6</sup> In addition, the chiral biimidazole also exhibited great potentials in asymmetric transformations. Kitamura and co-workers disclosed the dihydropyrroloimidazole-type bidentate chiral ligands (Naph-diPIM) from 2,7-dibromo-1,8-dinitronaphthalene and appropriate acetoneides of optically active 3,4-dihydroxy-pyrrolidine-2-one which was found utility in asymmetric Ru-catalyzed dehydrative allylation and Cu-catalyzed Friedel–Crafts reaction of indoles with trifluoropyruvates with high enantioselectivity (Scheme 1b).<sup>7</sup>

Despite these significant achievements in chiral imidazole ligands, the reported chiral imidazole ligands are mainly concentrated on the central chiral ligands and their application is limited. Therefore, the design and synthesis of efficient and practical chiral imidazole ligands is highly desirable. Among the various types of imidazoles, 2,2'-biimidazoles served as the


Received: February 22, 2019

Published: March 28, 2019

bimeric analogues of imidazole have been used as ligands applying to copper-catalyzed reactions.<sup>8</sup> However, the design of 2,2'-biimidazole compounds as chiral ligands has rarely been reported. These difficulties can be attributed to the lack of efficient method introducing chiral elements into the backbone of 2,2'-biimidazoles. Notably, the strategy of remote chirality delivery plays an important role in the design of axially chiral ligands, demonstrated by the elegant case of C3-TunePhos and their derivatives<sup>9</sup> and axially chiral 2,2'-bipyridine ligands (Cn-ACBP).<sup>10</sup> In these ligands, the chirality can be transferred from central to axial. Inspired by previous work, we envisioned that this remote chirality delivery strategy can be applied to synthesis of chiral 2,2'-biimidazole ligands (Scheme 1c). Herein, a series of binaphthyl-derived axially chiral 2,2'-biimidazole ligands has been designed and synthesized through the strategy of remote chirality delivery (chirality transfer from axial chirality to axial chirality). 2,2'-Biimidazole ligands can be fine-tuned by substitutions on the 3,3'-position of binaphthyl moiety<sup>11</sup> and a flexible backbone. These ligands have been proven to be efficient for Cu- or Fe-catalyzed asymmetric insertion of  $\alpha$ -aryl- $\alpha$ -diazoacetates into N–H bond of carbazoles with up to 96% ee.

The synthetic route is shown in Scheme 2, the treatment of (S)-3,3'-substituted-2,2'-bis(bromomethyl)-1,1'-binaphthalene

### Scheme 2. Syntheses of Axially Chiral 2,2'-Biimidazole Ligands



**1**<sup>12</sup> with 2,2'-biimidazoles **2** under basic conditions afforded a series of axially chiral 2,2'-biimidazole ligands **3** with moderate yields. This synthetic method features easy operation and provides diverse axially chiral 2,2'-biimidazole ligands in one step, which made it more potential in the application of asymmetric catalysis. The structure and absolute configuration of (S,S)-**3b** were determined by X-ray diffraction (XRD) analysis.<sup>13</sup> Notably, perfect remote chirality delivery could be realized through chirality transfer of binaphthyl moiety to 2,2'-biimidazole moiety.

With chiral 2,2'-biimidazole ligands in hand, we next started to investigate their performance in transition-metal-catalyzed asymmetric reactions. The chiral *N*-substituted carbazoles are core structure of numerous designed pharmaceutical molecules, which can serve as 5-HT receptor agonists for treatment of central nervous system diseases.<sup>14</sup> In 2017, Van Vranken and co-workers demonstrated Pd-catalyzed N–H insertion reaction between carbazoles and  $\alpha$ -diazo- $\alpha$ -phenylacetates by using a PyBOX, providing the products with excellent results.<sup>15</sup> However, high costs and toxicity of Pd reagents may restrict their wide application. As an abundant and inexpensive

transition metal, copper has been widely used as a metal precursor for catalytic asymmetric insertion of  $\alpha$ -diazoesters into N–H bonds.<sup>16</sup> To our delight, the copper complexes of (S,S)-**3a** was successfully applied in N–H insertion reaction between carbazoles and  $\alpha$ -aryl- $\alpha$ -diazoacetates, giving the desired products in 90% yield and 85% ee (see Table 1, entry 1).

Table 1. Evaluation of Reaction Parameters<sup>a</sup>

entry	solvent	temperature, T (°C)	(S,S)- <b>3</b>	yield <sup>b</sup> (%)	enantiomeric excess, ee <sup>c</sup> (%)
1	DCM	30	(S,S)- <b>3a</b>	90	85
2	CHCl <sub>3</sub>	30	(S,S)- <b>3a</b>	90	87
3	toluene (tol)	30	(S,S)- <b>3a</b>	92	86
4	THF	30	(S,S)- <b>3a</b>	72	7
5	<sup>t</sup> BuOMe	30	(S,S)- <b>3a</b>	74	92
6	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3a</b>	92	93
7	<sup>t</sup> BuOMe/tol (1:1)	40	(S,S)- <b>3a</b>	92	92
8	<sup>t</sup> BuOMe/tol (1:1)	50	(S,S)- <b>3a</b>	90	91
9	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3b</b>	82	81
10	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3c</b>	87	86
11	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3d</b>	82	90
12	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3e</b>	84	81
13	<sup>t</sup> BuOMe/tol (1:1)	30	(S,S)- <b>3f</b>	84	81

<sup>a</sup>Conditions: **4a** (0.10 mmol), **5a** (0.15 mmol), CuI (5.0 mol %), (S,S)-**3** (5.0 mol %), NaBAR<sup>F</sup> (12 mol %), 5 Å MS (200 mg), solvent (2.0 mL), 12 h. <sup>b</sup>Isolated yields. <sup>c</sup>Determined by chiral HPLC.

Original experiments employed  $\alpha$ -diazo- $\alpha$ -phenylacetate **4a** and carbazole **5a** at 30 °C in the presence of CuI/(S,S)-**3** as a catalyst and NaBAR<sup>F</sup> as an additive (Table 1). Initially, the reaction proceeded smoothly in DCM, giving the desired product **6aa** in 90% yield and 85% ee. Other different solvents were also evaluated, and it was found that the high yields were also achieved in toluene and chloroform, albeit with the moderate enantioselectivities. When the reaction solvent was THF, the yield was decreased to 72% (Table 1, entry 4). It was noted that *tert*-butyl methyl ether was proved to be beneficial for enantio-control, albeit comparatively low activity was observed (Table 1, entry 5). Fortunately, when a mixture solvent of toluene and <sup>t</sup>BuOMe with ratio of 1:1 was used, the optimal result was obtained (Table 1, entry 6). To further improve the efficiency of reaction, the temperature was increased to 40 and 50 °C, and yield was not obviously increased, although enantioselectivity slightly decreased. Therefore, we still selected 30 °C as the optimal temperature. Intrigued by these promising results, a series of axially chiral

2,2'-biimidazole ligands (*S,S*)-**3** were explored (Table 1, entries 9–13). We are pleased to find that all the chiral biimidazole ligands could give good results. The 2,2'-biimidazole ligands (*S,S*)-**3a** was proved to be beneficial for the reaction, furnishing the desired product in 93% ee and 92% yield. Thus, the optimized conditions were established as CuI (5.0 mol %)/(*S,S*)-**3a** (5.0 mol %)/NaBAR<sup>F</sup> (12 mol %)/5 Å MS/<sup>t</sup>BuOMe:Tol (1:1)/30 °C.

With the optimized reaction conditions established, we next investigated the substrate scope by reacting various  $\alpha$ -diazo- $\alpha$ -arylacetates **4** with carbazole **5a** (see Table 2). As expected,

**Table 2.** Cu-Catalyzed Asymmetric Insertion of  $\alpha$ -Aryl- $\alpha$ -diazoacetates into the N–H Bonds of Carbazoles<sup>a</sup>


entry	Ar	R	yield <sup>b</sup> (%)	enantiomeric excess, ee <sup>c</sup> (%)
1	Ph	Bn	92 (6aa)	93
2	4-MeC <sub>6</sub> H <sub>4</sub>	Bn	94 (6ba)	91
3	3-MeC <sub>6</sub> H <sub>4</sub>	Bn	91 (6ca)	93
4	4-MeOC <sub>6</sub> H <sub>4</sub>	Bn	88 (6da)	87
5	3-MeOC <sub>6</sub> H <sub>4</sub>	Bn	85 (6ea)	91
6	4-PhC <sub>6</sub> H <sub>4</sub>	Bn	94 (6fa)	90
7	4-FC <sub>6</sub> H <sub>4</sub>	Bn	93 (6ga)	92
8	4-ClC <sub>6</sub> H <sub>4</sub>	Bn	87 (6ha)	90
9	4-BrC <sub>6</sub> H <sub>4</sub>	Bn	89 (6ia)	89
10	3-BrC <sub>6</sub> H <sub>4</sub>	Bn	87 (6ja)	84
11	2-naphthyl	Bn	86 (6ka)	94
12	3-thienyl	Bn	81 (6la)	81
13	Ph	Me	86 (6ma)	88
14	Ph	Et	91 (6na)	90
15	Ph	<sup>t</sup> Pr	90 (6oa)	95

<sup>a</sup>Conditions: **4** (0.10 mmol), **5a** (0.15 mmol), CuI (5.0 mol %), (*S,S*)-**3a** (5.0 mol %), NaBAR<sup>F</sup> (12 mol %), 5 Å MS (200 mg), <sup>t</sup>BuOMe/tol (2.0 mL, 1:1), 12–24 h. <sup>b</sup>Isolated yields. <sup>c</sup>Determined by chiral HPLC.

various diazo substrates **4** performed very well under the optimized conditions. The influence of the steric and electronic properties of substituent on the phenyl ring of **4** was exploited. The reaction gave the desired products with high yields and enantioselectivities, when electron-donating groups were introduced into the phenyl ring of **4**. Notably, the steric effects of the aryl substituent of **4** rarely influenced the reaction (**6aa**–**6ea**). Substrates **4** with electron-withdrawing groups on the phenyl ring had little effect on enantioselectivity. Although electron-withdrawing groups obviously slowed the reaction down, satisfactory yields were still achieved by prolonging the reaction time (**6ga**–**6ja**). In the case of 2-naphthyl-substituted diazoacetates, the reaction gave the corresponding product with 94% ee and 86% yield (**6ka**). Furthermore, heteroaromatic thienyl-substituted diazoacetates also were suitable, affording the desired product with 81% ee and 81% yield (**6la**). Finally, the reaction of diazoacetates bearing different ester groups was examined, and a slightly higher enantioselectivity was observed upon increasing the steric hindrance (**6ma**–**6oa**). The isopropyl group afforded a higher ee value (95% ee).

To further demonstrate the versatility of our method. The second stage of exploration of the reaction scope focused on various substituted carbazoles (Scheme 3). Pleasingly, good

**Scheme 3.** Cu-Catalyzed Asymmetric Insertion of  $\alpha$ -Aryl- $\alpha$ -diazoacetates into the N–H Bonds of Carbazoles<sup>a</sup>



<sup>a</sup>Conditions: **4o** (0.10 mmol), **5** (0.15 mmol), CuI (5.0 mol %), (*S,S*)-**3a** (5.0 mol %), NaBAR<sup>F</sup> (12 mol %), 5 Å MS (200 mg), <sup>t</sup>BuOMe/tol (2.0 mL, 1:1), 24 h.


yields and excellent enantioselectivities were also achieved under the standard conditions. The results indicated that this protocol was compatible with electron-donating group as well as electron-withdrawing groups at the 2-position of carbazoles (**6ob**–**6od**). For the 3-methyl-9H-carbazole (**5e**), the reaction proceeded smoothly, delivering the product **6oe** with 84% yield and 94% ee. In addition, the carbazoles with a methyl group or a bromo group at the 4-position were also converted to the target products with excellent enantioselectivities (**6of**–**6og**).

The initial success of our chiral 2,2'-biimidazole ligands prompted us to further evaluate their practical utility. Iron, as an abundant and environmentally benign metal, is an ideal alternative to the precious metals. Iron catalysts could be used in the asymmetric transformations of diazo compounds in view of the facile change of oxidation state and the distinct Lewis acid character.<sup>17</sup> To best of our knowledge, the enantioselective catalytic N–H insertion reaction by using iron complexes as catalyst remains a great challenge to organic chemists.<sup>18</sup> We decided to further evaluate the above 2,2'-biimidazole ligands in iron-catalyzed enantioselective insertion of the N–H bonds of carbazoles.

Initially, the optimized reaction condition was established by a series of screening experiments. Using Fe(OTf)<sub>2</sub>/*(S,S)*-**3a** as a catalyst in *p*-xylene at 30 °C and NaBAR<sup>F</sup> as an additive, the reaction was conducted smoothly to give the target product in 72% yield and 93% ee (see Table S1 in the Supporting Information for details). All the chiral 2,2'-biimidazole ligands were evaluated, it was found that the electron-deficient ligands ((*S,S*)-**3c**, (*S,S*)-**3e**) gave higher yields and enantioselectivities than electron-rich ligands ((*S,S*)-**3b**, (*S,S*)-**3d**, (*S,S*)-**3f**). When (*S,S*)-**3a** was used, the highest ee value (93%) was obtained.

Under the optimized reaction conditions, several substrates were briefly examined for N–H insertion. As shown in Scheme 4, all the reaction could conduct smoothly, giving the products

**Scheme 4. Fe-Catalyzed Asymmetric Insertion of  $\alpha$ -Aryl- $\alpha$ -diazoacetates into the N–H Bonds of Carbazoles<sup>a</sup>**



<sup>a</sup>Conditions: **4** (0.10 mmol), **5** (0.15 mmol), Fe(OTf)<sub>2</sub> (5.0 mol %), (S,S)-**3a** (5.0 mol %), NaBARF (12 mol %), *p*-xylene (2.0 mL), 24 h.

in moderate yields and high enantioselectivities. First, the influence of ester size of the diazo compounds on activity and enantioselectivity of the N–H insertion was investigated, the benzyl ester afforded product with higher enantioselectivity (**6aa**–**6oa**). The best enantioselectivity was obtained for 2-diazo-2-(naphthalen-2-yl)acetate (**6ka**). When the electron-withdrawing group (–Cl) was introduced into the phenyl ring of **4**, the yield was decreased to 63% (**6ha**). The methyl on carbazole substrate was tolerated (**6oe**), but lower enantioselectivity was obtained, possibly because of the hindrance of methyl group.

In summary, we have designed and developed a series of novel axially chiral 2,2'-biimidazole ligands from (S)-2,2'-bis(bromomethyl)-1,1'-binaphthalene and 2,2'-biimidazoles in one step through the strategy of remote chirality delivery, which have shown excellent enantioselectivities and activities for Cu-catalyzed asymmetric insertion of  $\alpha$ -aryl- $\alpha$ -diazoacetates into the N–H bond of carbazoles. Moreover, the highly enantioselective iron-catalyzed insertion of  $\alpha$ -diazoacetates into the N–H bond have been realized by using iron complexes of chiral 2,2'-biimidazole ligands as a catalyst, with up to 94% ee. Note that this study paves a new way for further development of chiral imidazole ligands and realizes the application of 2,2'-biimidazole ligands in asymmetric catalysis. Further applications of these chiral 2,2'-biimidazole ligands in other catalytic asymmetric reactions are under investigation and will be reported in due course.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b00687.

General procedures and NMR spectra of obtained compounds (PDF)

## Accession Codes

CCDC 1887410 and 1887411 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [ygzhou@dicp.ac.cn](mailto:ygzhou@dicp.ac.cn).

### ORCID

Yong-Gui Zhou: 0000-0002-3321-5521

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

Financial support from the National Natural Science Foundation of China (Nos. 21532006, 21690074) and Dalian Institute of Chemical Physics (No. DMTO201501), and the Chinese Academy of Sciences (Nos. XDB17020300, QYZDJ-SSW-SLH035) is acknowledged.

## ■ REFERENCES

- (1) (a) Pozharskii, A. F.; Katritzky, A. R.; Soldatenkov, A. T. *Heterocycles in Life and Society: An Introduction to Heterocyclic Chemistry, Biochemistry and Applications*; Wiley: Chichester, U.K., 2011. (b) Fox, S. W. Chemistry of the Biologically Important Imidazoles. *Chem. Rev.* **1943**, 32, 47. (c) Sundberg, R. J.; Martin, R. B. Interactions of Histidine and Other Imidazole Derivatives with Transition Metal Ions in Chemical and Biological Systems. *Chem. Rev.* **1974**, 74, 471. (d) Ingle, R. A. Histidine Biosynthesis. In *The Arabidopsis Book*, Vol. 9; American Society of Plant Biologists: Rockville, MD, 2011. (e) Kleingardner, J. G.; Bren, K. L. Biological Significance and Applications of Heme *c* Proteins and Peptides. *Acc. Chem. Res.* **2015**, 48, 1845.
- (2) For selected examples on imidazole-based organocatalysts, see: (a) Davie, E. A. C.; Mennen, S. M.; Xu, Y.; Miller, S. J. Asymmetric Catalysis Mediated by Synthetic Peptides. *Chem. Rev.* **2007**, 107, 5759. (b) Zhao, Y.; Mitra, A. W.; Hoveyda, A. H.; Snapper, M. L. Kinetic Resolution of 1,2-Diols through Highly Site- and Enantioselective Catalytic Silylation. *Angew. Chem., Int. Ed.* **2007**, 46, 8471. (c) Zhang, Z.; Xie, F.; Jia, J.; Zhang, W. Chiral Bicycle Imidazole Nucleophilic Catalysts: Rational Design, Facile Synthesis, and Successful Application in Asymmetric Steglich Rearrangement. *J. Am. Chem. Soc.* **2010**, 132, 15939. (d) Zhao, Y.; Rodrigo, J.; Hoveyda, A. H.; Snapper, M. L. Enantioselective Silyl Protection of Alcohols Catalysed by an Amino-Acid-Based Small Molecule. *Nature* **2006**, 443, 67.
- (3) For selected examples on imidazole-based ligands, see: (a) Diaconescu, P. L. Reactions of Aromatic *N*-Heterocycles with d<sup>9</sup>Pt-Metal Alkyl Complexes Supported by Chelating Diamide Ligands. *Acc. Chem. Res.* **2010**, 43, 1352. (b) Díez, J.; Gimeno, J.; Lledós, A.; Suárez, F. J.; Vicent, C. Imidazole Based Ruthenium(IV) Complexes as Highly Efficient Bifunctional Catalysts for the Redox Isomerization of Allylic Alcohols in Aqueous Medium: Water as Cooperating Ligand. *ACS Catal.* **2012**, 2, 2087. (c) Chen, S.-S. The Roles of Imidazole Ligands in Coordination Supramolecular Systems. *CrystEngComm* **2016**, 18, 6543. (d) Asano, K.; Matsubara, S. Effects of a Flexible Alkyl Chain on a Ligand for CuAAC Reaction. *Org. Lett.* **2010**, 12, 4988. (e) Song, B.; Knauber, T.; Gooßen, L. J. Decarboxylative Cross-Coupling of Mesylates Catalyzed by Copper/

Palladium Systems with Customized Imidazolyl Phosphine Ligands. *Angew. Chem., Int. Ed.* **2013**, *52*, 2954. (f) Nakatake, D.; Yokote, Y.; Matsushima, Y.; Yazaki, R.; Ohshima, T. A Highly Stable but Highly Reactive Zinc Catalyst for Transesterification Supported by a Bis(imidazole) Ligand. *Green Chem.* **2016**, *18*, 1524. (g) Grotjahn, D. B.; Kraus, J. E.; Amouri, H.; Rager, M.-N.; Cooksy, A. L.; Arita, A. J.; Cortes-Llamas, S. A.; Mallari, A. A.; DiPasquale, A. G.; Moore, C. E.; Liable-Sands, L. M.; Golen, J. D.; Zakharov, L. N.; Rheingold, A. L. Multimodal Study of Secondary Interactions in Cp\*Ir Complexes of Imidazolylphosphines Bearing an NH Group. *J. Am. Chem. Soc.* **2010**, *132*, 7919.

(4) For selected examples on imidazole-based chiral ligands, see: (a) Orlandi, S.; Benaglia, M.; Cozzi, F. Cu(II)-Catalyzed Enantioselective Aldol Condensation Between Malonic Acid Hemithioesters and Aldehydes. *Tetrahedron Lett.* **2004**, *45*, 1747. (b) Li, Y.; Ding, K.; Sandoval, C. A. Hybrid NH<sub>2</sub>-Benzimidazole Ligands for Efficient Ru-Catalyzed Asymmetric Hydrogenation of Aryl Ketones. *Org. Lett.* **2009**, *11*, 907. (c) Zhang, Z.-G.; Dong, Z.-B.; Li, J.-S. Synthesis and Application of 3-Substituted (S)-BINOL as Chiral Ligands for the Asymmetric Ethylation of Aldehydes. *Chirality* **2010**, *22*, 820. (d) Li, X.-N.; Zhou, H.-Y.; Feng, L.; Duan, K.; Wang, J.-X. Temperature-tuned Enantioselectivity in Transfer Hydrogenation: Preparation and Catalytic Property of Amino Acid-modified Benzimidazole. *Appl. Organomet. Chem.* **2012**, *26*, 168. (e) Zhou, Y.; Zhu, F.-L.; Liu, Z.-T.; Zhou, X.-M.; Hu, X.-P. Chiral Ferrocenyl P,N-Ligands for Palladium-Catalyzed Asymmetric Formal [3 + 2] Cycloaddition of Propargylic Esters with  $\beta$ -Ketoesters: Access to Functionalized Chiral 2,3-Dihydrofurans. *Org. Lett.* **2016**, *18*, 2734. (f) Quan, X.; Liu, J.; Rabten, W.; Diomed, S.; Singh, T.; Andersson, P. G. Thiazole, Imidazole and Oxazoline Based N,P-Ligands for Palladium-Catalyzed Cycloisomerization of 1,6-Enynes. *Eur. J. Org. Chem.* **2016**, *2016*, 3427. (g) Xu, D.; Sun, Q.; Quan, Z.; Sun, W.; Wang, X. The Synthesis of Chiral Tridentate Ligands from L-Proline and Their Application in the Copper(II)-catalyzed Enantioselective Henry Reaction. *Tetrahedron: Asymmetry* **2017**, *28*, 954.

(5) (a) Kaukoranta, P.; Engman, M.; Hedberg, C.; Bergquist, J.; Andersson, P. G. Iridium Catalysts with Chiral Imidazole-Phosphine Ligands for Asymmetric Hydrogenation of Vinyl Fluorides and other Olefins. *Adv. Synth. Catal.* **2008**, *350*, 1168. (b) Mazuela, J.; Tolstoy, P.; Pàmies, O.; Andersson, P. G.; Diéguez, M. Phosphite-oxazole/imidazole Ligands in Asymmetric Intermolecular Heck Reaction. *Org. Biomol. Chem.* **2011**, *9*, 941. (c) Paptchikhine, A.; Itto, K.; Andersson, P. G. Sequential Birch Reaction and Asymmetric Ir-Catalyzed Hydrogenation as a Route to Chiral Building Blocks. *Chem. Commun.* **2011**, *47*, 3989. (d) Li, J.-Q.; Liu, J.; Krajangrsi, S.; Chumanvej, N.; Singh, T.; Andersson, P. G. Asymmetric Hydrogenation of Allylic Alcohols Using Ir-N,P-Complexes. *ACS Catal.* **2016**, *6*, 8342.

(6) (a) Cardoso, F. S. P.; Abboud, K. A.; Aponick, A. Design, Preparation, and Implementation of an Imidazole-Based Chiral Biaryl P,N-Ligand for Asymmetric Catalysis. *J. Am. Chem. Soc.* **2013**, *135*, 14548. (b) Pappoppula, M.; Cardoso, F. S. P.; Garrett, B. O.; Aponick, A. Enantioselective Copper-Catalyzed Quinoline Alkynylation. *Angew. Chem., Int. Ed.* **2015**, *54*, 15202. (c) Paioti, P. H. S.; Abboud, K. A.; Aponick, A. Catalytic Enantioselective Synthesis of Amino Skipped Dienes. *J. Am. Chem. Soc.* **2016**, *138*, 2150. (d) Mishra, S.; Liu, J.; Aponick, A. Enantioselective Alkyne Conjugate Addition Enabled by Readily Tuned Atropisomeric P,N-Ligands. *J. Am. Chem. Soc.* **2017**, *139*, 3352.

(7) (a) Miyata, K.; Kutsuna, H.; Kawakami, S.; Kitamura, M. A Chiral Bidentate sp<sup>2</sup>-N Ligand, Naph-diPIM: Application to CpRu-Catalyzed Asymmetric Dehydrative C-, N-, and O-Allylation. *Angew. Chem., Int. Ed.* **2011**, *50*, 4649. (b) Kitamura, M.; Miyata, K.; Seki, T.; Vatmurge, N.; Tanaka, S. CpRu-Catalyzed Asymmetric Dehydrative Allylation. *Pure Appl. Chem.* **2013**, *85*, 1121. (c) Suzuki, Y.; Vatmurge, N.; Tanaka, S.; Kitamura, M. Enantio- and Diastereo-selective Dehydrative "One-Step" Construction of Spirocarbocycles via a Ru/H<sup>+</sup>-Catalyzed Tsuji-Trost Approach. *Chem. - Asian J.* **2017**, *12*, 633. (d) Le, T. P.; Higashita, K.; Tanaka, S.; Yoshimura, M.;

Kitamura, M. Revisiting the Cu<sup>II</sup>-Catalyzed Asymmetric Friedel-Crafts Reaction of Indole with Trifluoropyruvate. *Org. Lett.* **2018**, *20*, 7149.

(8) (a) Wang, B.-A.; Zeng, R.-S.; Wei, H.-Q.; Jia, A.-Q.; Zou, J.-P. Copper(II) and 2,2'-Biimidazolyl-promoted Ullmann Coupling Reaction of Phenols and Aryl Iodides. *Chin. J. Chem.* **2006**, *24*, 1062. (b) Hu, Z.; Ye, W.; Zou, H.; Yu, Y. 2,2'-Biimidazole as an Efficient Ligand for Copper(I)-Catalyzed C-N Coupling Reactions. *Synth. Commun.* **2009**, *40*, 222. (c) Zong, C.; Liu, J.; Chen, S.; Zeng, R.; Zou, J.-P. Efficient C-S Cross-Coupling of Thiols with Aryl Iodides Catalyzed by Cu(OAc)<sub>2</sub>·H<sub>2</sub>O and 2,2'-Biimidazole. *Chin. J. Chem.* **2014**, *32*, 212. (d) Hu, Z.; Ou, L.-L.; Li, S.-D.; Yang, L. Efficient Copper(I)-Catalyzed, Microwave-assisted, One-pot Synthesis of 3,4-Diaryl Isoquinolines. *Res. Chem. Intermed.* **2015**, *41*, 3461. (e) Nitelet, A.; Wouters, J.; Dewez, D. F.; Evano, G. Copper-Catalyzed Alkenylation of Cyanamides. *Org. Lett.* **2017**, *19*, 6276.

(9) (a) Qiu, L.; Kwong, F. Y.; Wu, J.; Lam, W. H.; Chan, S.; Yu, W.-Y.; Li, Y.-M.; Guo, R.; Zhou, Z.; Chan, A. S. C. A New Class of Versatile Chiral-Bridged Atropisomeric Diphosphine Ligands: Remarkably Efficient Ligand Syntheses and Their Applications in Highly Enantioselective Hydrogenation Reactions. *J. Am. Chem. Soc.* **2006**, *128*, 5955. (b) Sun, X.; Zhou, L.; Li, W.; Zhang, X. Convenient Divergent Strategy for the Synthesis of TunePhos-Type Chiral Diphosphine Ligands and Their Applications in Highly Enantioselective Ru-Catalyzed Hydrogenations. *J. Org. Chem.* **2008**, *73*, 1143.

(10) Gao, X.; Wu, B.; Huang, W.-X.; Chen, M.-W.; Zhou, Y.-G. Enantioselective Palladium-Catalyzed C-H Functionalization of Indoles Using an Axially Chiral 2,2'-Bipyridine Ligand. *Angew. Chem., Int. Ed.* **2015**, *54*, 11956.

(11) (a) Brunel, J. M. BINOL: A Versatile Chiral Reagent. *Chem. Rev.* **2005**, *105*, 857. (b) Noyori, R. Asymmetric Catalysis: Science and Opportunities (Nobel Lecture). *Angew. Chem., Int. Ed.* **2002**, *41*, 2008.

(12) Ooi, T.; Kameda, M.; Maruoka, K. Design of N-Spiro C<sub>2</sub>-Symmetric Chiral Quaternary Ammonium Bromides as Novel Chiral Phase-Transfer Catalysts: Synthesis and Application to Practical Asymmetric Synthesis of  $\alpha$ -Amino Acids. *J. Am. Chem. Soc.* **2003**, *125*, 5139.

(13) (S,S)-**3b** and palladium complex [Pd(3c)Cl<sub>2</sub>·(CH<sub>3</sub>COCH<sub>3</sub>)<sub>2</sub>] were confirmed by X-ray crystallography.

(14) Tenbrink, R. E. Oxazinocarbazoles for the Treatment of CNS Diseases, U.S. Patent No. US 6821970(B2), Nov. 23, 2004.

(15) Arredondo, V.; Hiew, S. C.; Gutman, E. S.; Premachandra, I. D. U. A.; Van Vranken, D. L. Enantioselective Palladium-Catalyzed Carbene Insertion into the N-H Bonds of Aromatic Heterocycles. *Angew. Chem., Int. Ed.* **2017**, *56*, 4156.

(16) For selected reviews and examples on Cu-catalyzed insertion of  $\alpha$ -diazoesters into N-H bonds, see: (a) Zhu, S.-F.; Zhou, Q.-L. Transition-Metal-Catalyzed Enantioselective Heteroatom-Hydrogen Bond Insertion Reactions. *Acc. Chem. Res.* **2012**, *45*, 1365. (b) Bachmann, S.; Fielenbach, D.; Jørgensen, K. A. Cu(I)-Carbenoid- and Ag(I)-Lewis Acid-Catalyzed Asymmetric Intermolecular Insertion of  $\alpha$ -Diazo Compounds into N-H Bonds. *Org. Biomol. Chem.* **2004**, *2*, 3044. (c) Lee, E. C.; Fu, G. C. Copper-Catalyzed Asymmetric N-H Insertion Reactions: Couplings of Diazo Compounds with Carbamates to Generate  $\alpha$ -Amino Acids. *J. Am. Chem. Soc.* **2007**, *129*, 12066. (d) Liu, B.; Zhu, S.-F.; Zhang, W.; Chen, C.; Zhou, Q.-L. Highly Enantioselective Insertion of Carbenoids into N-H Bonds Catalyzed by Copper Complexes of Chiral Spiro Bisoaxazolines. *J. Am. Chem. Soc.* **2007**, *129*, 5834. (e) Hou, Z.; Wang, J.; He, P.; Wang, J.; Qin, B.; Liu, X.; Lin, L.; Feng, X. Highly Enantioselective Insertion of Carbenoids into N-H Bonds Catalyzed by Copper(I) Complexes of Binol Derivatives. *Angew. Chem., Int. Ed.* **2010**, *49*, 4763. (f) Zhu, S.-F.; Xu, B.; Wang, G.-P.; Zhou, Q.-L. Well-Defined Binuclear Chiral Spiro Copper Catalysts for Enantioselective N-H Insertion. *J. Am. Chem. Soc.* **2012**, *134*, 436.

(17) For selected reviews and examples on Fe-catalyzed asymmetric transformations of diazo compounds, see: (a) Sun, X.-L.; Zheng, J.-C.; Tang, Y. Iron Carbenoid-mediated Ylide Reactions. *Pure Appl. Chem.* **2010**, *82*, 625. (b) Gopalaiah, K. Chiral Iron Catalysts for Asymmetric

Synthesis. *Chem. Rev.* **2013**, *113*, 3248. (c) Zhu, S.-F.; Zhou, Q.-L. Iron-Catalyzed Transformations of Diazo Compounds. *Natl. Sci. Rev.* **2014**, *1*, 580. (d) Coelho, P. S.; Brustad, E. M.; Kannan, A.; Arnold, F. H. Olefin Cyclopropanation *via* Carbene Transfer Catalyzed by Engineered Cytochrome P450 Enzymes. *Science* **2013**, *339*, 307. (e) Li, C.-Y.; Wang, X.-B.; Sun, X.-L.; Tang, Y.; Zheng, J.-C.; Xu, Z.-H.; Zhou, Y.-G.; Dai, L.-X. Iron Porphyrin-Catalyzed Olefination of Ketenes with Diazoacetate for the Enantioselective Synthesis of Allenes. *J. Am. Chem. Soc.* **2007**, *129*, 1494. (f) Zhu, S.-F.; Cai, Y.; Mao, H.-X.; Xie, J.-H.; Zhou, Q.-L. Enantioselective Iron-Catalyzed O-H Bond Insertions. *Nat. Chem.* **2010**, *2*, 546. (g) Xu, H.; Li, Y.-P.; Cai, Y.; Wang, G.-P.; Zhu, S.-F.; Zhou, Q.-L. Highly Enantioselective Copper- and Iron-Catalyzed Intramolecular Cyclo-propanation of Indoles. *J. Am. Chem. Soc.* **2017**, *139*, 7697.

(18) (a) Aviv, I.; Gross, Z. Iron Corroles and Porphyrins as Very Efficient and Highly Selective Catalysts for the Reactions of  $\alpha$ -Diazo Esters with Amines. *Synlett* **2006**, *2006*, 951. (b) Baumann, L. K.; Mbuvi, H. M.; Du, G.; Woo, L. K. Iron Porphyrin Catalyzed N-H Insertion Reactions with Ethyl Diazoacetate. *Organometallics* **2007**, *26*, 3995. (c) Mbuvi, H. M.; Klobukowski, E. R.; Roberts, G. M.; Woo, L. K. O-H Insertion and Tandem N-H Insertion/cyclization Reactions Using an Iron Porphyrin as Catalyst with Diazo Compounds as Carbene Sources. *J. Porphyrins Phthalocyanines* **2010**, *14*, 284.